XXXIII International (ONLINE) Workshop on High Energy Physics "Hard Problems of Hadron Physics: Non-Perturbative QCD & Related Quests"

Contribution ID: 36

Type : not specified

Jet transport coefficient qhat in lattice QCD

пятница, 12 ноября 2021 г. 12:00 (30)

We present the first calculation of the jet transport coefficient \hat{q} in quenched and (2+1)-flavor QCD on a 4-D Euclidean lattice. The light-like propagation of an energetic parton is factorized from the mean square gain in momentum transverse to the direction of propagation, which is expressed in terms of the thermal field-strength field-strength correlator. The leading-twist term in its operator product expansion is calculated on the lattice. Continuum extrapolated quenched results, and full QCD estimates based on un-renormalized lattice data, over multiple lattice sizes, are compared with (non) perturbative calculations and phenomenological extractions of \hat{q} . The lattice data for \hat{q} show a temperature dependence similar to the entropy density. Within uncertainties, these are consistent with phenomenological extractions, contrary to calculations using perturbation theory.

Primary author(s) : WEBER, Johannes Heinrich (Humboldt-Universität zu Berlin)

Co-author(s) : Dr KUMAR, Amit (McGill University); Prof. MAJUMDER, Abhijit (Wayne State University)

Presenter(s): WEBER, Johannes Heinrich (Humboldt-Universität zu Berlin)

Session Classification : Morning session 9

Track Classification : Quark-Gluon Matter at Finite Densities and Temperatures